

INTERNATIONAL UNION OF PURE
AND APPLIED CHEMISTRY

PHYSICAL CHEMISTRY DIVISION
COMMISSION ON MOLECULAR STRUCTURE AND SPECTROSCOPY*

PARAMETERS AND SYMBOLS FOR USE IN
NUCLEAR MAGNETIC RESONANCE

(IUPAC Recommendations 1997)

Prepared for publication by

ROBIN K. HARRIS¹, JOZEF KOWALEWSKI² AND SONIA CABRAL DE MENEZES³

¹Department of Chemistry, University of Durham, South Road, Durham DH1 3LE, UK.

²Division of Physical Chemistry, Arrhenius Laboratory, Stockholm University, S-10691 Stockholm, Sweden.

³PETROBRAS/CENPES, Divisão de Química, Ilha do Fundão, Quadra 7, Cidade Universitaria, 21949 900 Rio de Janeiro, R.J., Brazil

*Membership of the Commission during the preparation of this report (1995–97) was as follows:

Chairman: J. E. Bertie (Canada; 1994–97); **Secretary:** P. Klæboe (Norway; 1996–97); **Titular Members:** A. M. Heyns (RSA; 1994–97); R. Janoschek (Austria; 1994–97); P. Klæboe (Norway; 1994–97); S. Tsuchiya (Japan; 1994–97); B. P. Winnewisser (Germany; 1991–97); **Associate Members:** A. M. Bradshaw (FRG; 1991–95); S. M. Cabral de Menezes (Brazil; 1994–97); B. G. Derendjaev (Russia; 1991–95); E. Hirota (Japan; 1994–97); J. Kowalewski (Sweden; 1996–97); A. Oskam (Netherlands; 1991–97); C. Zhang (China; 1989–97); **National Representatives:** J. E. Collin (Belgium; 1985–97); M. Chowdhury (India; 1986–95); S. Califano (Italy; 1989–97); J. J. C. Teixeira-Dias (Portugal; 1991–97); Y. S. Lee (Republic of Korea; 1989–97); D. Escolar (Spain; 1991–97); J. Kowalewski (Sweden; 1994–95); S. Suzer (Turkey; 1996–97); S. Içli (Turkey; 1994–95); R. K. Harris (UK; 1994–97).

Republication or reproduction of this report or its storage and/or dissemination by electronic means is permitted without the need for formal IUPAC permission on condition that an acknowledgement, with full reference to the source along with use of the copyright symbol ©, the name IUPAC and the year of publication are prominently visible. Publication of a translation into another language is subject to the additional condition of prior approval from the relevant IUPAC National Adhering Organization.

Parameters and symbols for use in nuclear magnetic resonance (IUPAC Recommendations 1997)

Abstract: NMR is now frequently the technique of choice for the determination of chemical structure in solution. Its uses also span structure in solids and mobility at the molecular level in all phases. The research literature in the subject is vast and ever-increasing. Unfortunately, many articles do not contain sufficient information for experiments to be repeated elsewhere, and there are many variations in the usage of symbols for the same physical quantity. It is the aim of the present recommendations to provide simple check-lists that will enable such problems to be minimised in a way that is consistent with general IUPAC formulation. The area of medical NMR and imaging is not specifically addressed in these recommendations, which are principally aimed at the mainstream use of NMR by chemists (of all sub-disciplines) and by many physicists, biologists, materials scientists and geologists etc. working with NMR.

The document presents recommended notation for use in journal publications involving a significant contribution of nuclear magnetic resonance (NMR) spectroscopy. The recommendations are in two parts:

A. Experimental parameters which should be listed so that the work in question can be repeated elsewhere.

B. A list of symbols (using Roman or Greek characters) to be used for quantities relevant to NMR.

A. NMR EXPERIMENTAL PARAMETERS for reporting in publications

1. Introduction

The aim of this list is to introduce a degree of uniformity in the way in which NMR results are reported in chemical journals, so as to ensure that experiments can be precisely repeated. It is to be regarded as a minimal list, though not all parameters are relevant to a given experiment. It should be noted that sample conditions (e.g. solvent, concentration etc.) are not part of the list, though clearly it is also essential for these to be specified.

2. General

- (a) The nucleus or nuclei studied.
- (b) The instrument manufacturer and model.
- (c) The resonance frequency or frequencies (preferably of the reference sample or samples).
- (d) The magnetic field [optional if frequencies are given].
- (e) The type(s) of pulse sequence.
- (f) The referencing procedure for obtaining chemical shifts (and/or relative shielding constants).
- (g) The size/nature of the sample tube/container.
- (h) The temperature of the sample at the time of the measurement.

3. Pulse conditions

- (a) The RF pulse magnetic field (or equivalent in frequency units) and/or equivalent 90° pulse duration. [Note the unambiguous word "duration" rather than "width" or "length".]
- (b) The flip (nutation) angle(s) and relative phases of all pulses.
- (c) The spectral width (or dwell time if unambiguously defined).
- (d) The acquisition time (or number of acquired data points if the dwell time is given).

- (e) The number of transients (acquisitions).
- (f) The recycle delay. [Note, this implies the time interval between the completion of one cycle of a pulse sequence and the start of the next one, sometimes called a relaxation delay. This is distinct from a cycle time, which is the time between analogous points of successive pulse sequence cycles].
- (g) Any other relevant time intervals.
- (h) The decoupling or double/multiple resonance method and conditions, including the RF magnetic field (or equivalent in frequency units and/or equivalent 90° pulse duration).
- (i) Pulse shapes and relevant frequencies (for experiments with selective pulses).
- (j) The magnitudes, directions, shapes and durations of any magnetic field gradients.

4. Processing parameters

- (a) Zero-filling extent.
- (b) Any window function (apodization, pre-acquisition delay, resolution enhancement, line broadening).
- (c) Any special baseline correction or phase adjustment procedures.
- (d) Any additional processing details, such as linear prediction or maximum entropy.
- (e) Any special procedures for signal quantification (e.g. deconvolution).

5. Solid-state spectra (additional requirements)

- (a) Contact time. [For cross polarization operation.]
- (b) Sample rotation angle.
- (c) Sample rotation rate.
- (d) Use of special techniques such as DOR or DAS.

6. Relaxation measurements (additional requirements)

- (a) Number of measurement points used.
- (b) Analysis method (exponential etc.).
- (c) Radiofrequency field magnitude (or equivalent expressed in frequency units). [For $T_{1\rho}$ measurements.]

7. Multidimensional spectra

- (a) The time increments and the number of data points and/or the spectral width and maximum duration of evolution periods must be given for each dimension, as well as the data size after processing.
- (b) For figures of multidimensional spectra, each axis should be unambiguously labelled by the variable involved (e.g. δ_{H} , δ_{C} , J_{CH}) where possible, as well as by a frequency label such as F_1 , F_2 etc.

8. Additional notes

- (a) Figures of spectra should have an indication of the direction of increasing frequency and the captions should state the nucleus involved. Wherever feasible, assignments of chemical shifts should be indicated on illustrated spectra.
- (b) Phase cycling of all RF pulses and of the receiver should be specified when crucial to the experiment.
- (c) For novel or unusual pulse sequences a timing diagram is highly recommended. If acronyms are used instead of diagrams for established sequences, a literature reference is recommended.

B. SYMBOLS FOR NMR AND RELATED QUANTITIES

(i) Roman alphabet

a or *A* Hyperfine (electron-nucleus) coupling constant

$A_q^{(l,m)}$	The m th component of an irreducible tensor of order l representing the nuclear spin operator for an interaction of type q
B	Magnetic field (strictly the magnetic flux density or magnetic induction)
B_0	Static magnetic field of an NMR spectrometer
B_1, B_2	Radiofrequency magnetic fields associated with frequencies ν_1, ν_2
B_L	Local magnetic field (components B_{xL}, B_{yL}, B_{zL}) of random field or dipolar origin
C	Spin-rotation interaction tensor
C_X	Spin-rotation coupling constant of nuclide X
D	Dipolar interaction tensor
D	Dipolar coupling constant between two nuclei (say 1 and 2), $(\mu_0/4\pi)\gamma_1\gamma_2(\hbar/2\pi)r_{12}^{-3}$, in frequency units (footnote 1).
D^C	Nuclear receptivity relative to that of the carbon-13 nucleus
D^P	Nuclear receptivity relative to that of the proton (hydrogen-1 nucleus)
E	Electric field strength
F	Spectral width
F_1, F_2 (or f_1, f_2)	The two frequency dimensions of a two-dimensional spectrum (use F_3 etc. for higher orders)
\hat{F}_G	Nuclear spin operator for a group, G , of nuclei (components $\hat{F}_{Gx}, \hat{F}_{Gy}, \hat{F}_{Gz}, \hat{F}_{G+}, \hat{F}_{G-}$)
F_G	Magnetic quantum number associated with \hat{F}_G
g	Nuclear or electronic g factor (Landé splitting factor)
G	Magnetic field gradient amplitude
H_{ij}	Element of matrix representation of Hamiltonian operator (in energy units); subscripts indicate the nature of the operator
\hat{I}_j	Nuclear spin operator for nucleus j (components $\hat{I}_{jx}, \hat{I}_{jy}, \hat{I}_{jz}$)
$\hat{I}_{j+}, \hat{I}_{j-}$	'Raising' and 'lowering' spin operators for nucleus j
I_j	Magnetic quantum number associated with \hat{I}_j
J	Indirect coupling tensor
nJ	Nuclear spin-spin coupling constant through n bonds (usually given in frequency units). Parentheses may be used (for example) to indicate the species of nuclei coupled, e.g. $J(^{13}\text{C}, ^1\text{H})$ or, additionally, the coupling path, e.g. $J(\text{POCF})$. Where no ambiguity arises, the elements involved can be, alternatively, given as subscripts, e.g. J_{CH} . The nucleus of higher mass should be given first
$J(\omega)$	Spectral density of fluctuations at angular frequency ω . Subscripts and superscripts to J may be used to indicate the relevant quantum number change (0, 1 or 2) or the order and component of the relevant tensor quantities.
nK	Reduced nuclear spin-spin coupling constant (see the notes concerning nJ), $K_{jk} = 4\pi^2 J_{jk}/h\gamma_j\gamma_k$
L	Angular momentum
m_j	Eigenvalue of \hat{I}_{jz} (magnetic component quantum number) (footnote 2)

m_{tot}	Total magnetic component number for a spin system (eigenvalue of $\sum_j \hat{I}_{jz}$) (footnote 2)
$m_{\text{tot}}(\text{X})$	Total magnetic component quantum number for X-type nuclei (footnote 2)
M_0	Equilibrium macroscopic magnetization per volume of a spin system in the presence of B_0
M_x, M_y, M_z	Components of macroscopic magnetization per volume.
M_n	n th moment of spectrum (M_2 = second moment, etc.)
n_α, n_β	Populations of the α and β spin states
N	Total number of nuclei of a given type per volume in the sample
\mathbf{q}	Electric field gradient tensor in units of the elementary charge (principal components q_{XX}, q_{YY}, q_{ZZ}) (see also \mathbf{V})
Q	eQ is the nuclear quadrupole moment, where e is the elementary charge
R_1^X	Spin-lattice (longitudinal) relaxation rate constant for nucleus X
R_2^X	Spin-spin (transverse) relaxation rate constant for nucleus X
$R_{1\rho}^X$	Spin-lattice relaxation rate constant in the rotating frame for nucleus X
S	Signal intensity
\hat{S}	Electron (or, occasionally, nuclear) spin operator; cf. \hat{I}
t_1, t_2	Time dimensions for two-dimensional NMR
T_c	Coalescence temperature for signals in an NMR spectrum
T_1^X	Spin-lattice (longitudinal) relaxation time of the X nucleus (further subscripts refer to the relaxation mechanism)
T_2^X	Spin-spin (transverse) relaxation time of the X nucleus (further subscripts refer to the relaxation mechanism)
T_2^*	Net dephasing time for M_x or M_y (including contribution from magnetic field inhomogeneity)
$T_{1\rho}^X$	Spin-lattice relaxation time of the X nucleus in the frame of reference rotating with B_1
T_d	Pulse (recycle) delay
T_{ac}	Acquisition time
$T_q^{(l,m)}$	The m th component of an irreducible tensor of order l representing the strength of an interaction of type q
\mathbf{V}	Electric field gradient tensor. $\mathbf{V} = e\mathbf{q}$, where e is the elementary charge
$V_{\alpha\beta}$	Elements of Cartesian electric field gradient tensor
W_0, W_1, W_2	Relaxation rate constants (transition probabilities per time) between energy levels differing by 0, 1, and 2 (respectively) in m_{tot} ; especially, but not uniquely, for systems of two spin- $\frac{1}{2}$ nuclei
W_{rs}	Transition probability between spin states r and s

(ii) Greek alphabet

α	Nuclear spin wavefunction (eigenfunction of \hat{I}_{jz}) for the $m_I = +\frac{1}{2}$ state of a spin- $\frac{1}{2}$ nucleus
α_E	The Ernst angle (for optimum sensitivity)

β	Nuclear spin wavefunction (eigenfunction of \hat{I}_z) for the $m_I = -\frac{1}{2}$ state of a spin- $\frac{1}{2}$ nucleus
γ_X	Magnetogyric ratio of nucleus X
δ_X	Chemical shift (for the resonance) of nucleus of element X (positive when the sample resonates to high frequency of the reference). Usually in ppm (footnote 3). Further information regarding solvent, references, or nucleus of interest may be given by superscripts or subscripts or in parentheses.
Δn	Population difference between nuclear states (Δn_0 at Boltzmann equilibrium)
$\Delta\delta$	Change or difference in δ
$\Delta\nu_{\frac{1}{2}}$	Full width in frequency units of a resonance line at half-height
$\Delta\sigma$	Anisotropy in σ [$\Delta\sigma = \sigma_{ZZ} - \frac{1}{3}(\sigma_{XX} + \sigma_{YY})$] (footnote 4). (see also ζ)
$\Delta\chi$	(i) Susceptibility anisotropy ($\Delta\chi = \chi_{ } - \chi_{\perp}$); (ii) difference in electronegativities
ϵ_0	Permittivity of a vacuum
ζ	Anisotropy in shielding (footnote 4), expressed as $\sigma_{ZZ} - \sigma_{\text{iso}}$. (see also $\Delta\sigma$)
η	(i) Nuclear Overhauser enhancement (so that the nuclear Overhauser effect is $1 + \eta$); (ii) tensor asymmetry factor (e.g. in σ); (iii) viscosity
κ	Skew of a tensor. (see also footnote 7)
θ	Angle, especially for that between a given vector and B_0
μ	(i) Magnetic dipole moment (component μ_z along B_0); (ii) electric dipole moment
μ_0	Permeability of a vacuum
μ_B	Bohr magneton
μ_N	Nuclear magneton
ν_j	Larmor precession frequency of nucleus j (usually given in MHz)
ν_0	(i) Spectrometer operating frequency; (ii) Larmor precession frequency (general, or of bare nucleus)
ν_1	Frequency of 'observing' RF magnetic field B_1 (to be distinguished from its strength, γB_1 , for which the symbol Ω_1 is recommended).
ν_2	Frequency of 'irradiating' RF magnetic field B_2 (to be distinguished from its strength, γB_2 , for which the symbol Ω_2 is recommended).
$\bar{\nu}_X$	Resonance frequency for the nucleus of element X in a magnetic field such that the protons in tetramethylsilane (TMS) resonate at exactly 100 MHz
ρ	Density matrix
$\hat{\rho}$	Density operator
ρ_{ij}	Element of matrix representation of $\hat{\rho}$
σ	Shielding tensor (footnotes 5 and 6)
σ_j	(Isotropic) shielding constant of nucleus j . Usually given in ppm. Subscripts may alternatively indicate contributions to σ
$\sigma_{ }, \sigma_{\perp}$	Components of shielding tensor σ parallel and perpendicular to the symmetry axis (axially-symmetric case) (footnote 5)

$\hat{\sigma}$	Reduced density operator
τ	(i) Time between RF pulses (general symbol); (ii) lifetime in dynamic NMR usage
τ_c	Correlation time for molecular-level motion, especially for isotropic molecular tumbling
τ_d	Dwell time
τ_{null}	Recovery time sufficing to give zero signal after a 180 ° pulse
τ_p	Pulse duration
τ_{sc}	Correlation time for relaxation by the scalar mechanism
τ_{sr}	Correlation time for spin-rotation relaxation
$\tau_{\parallel}, \tau_{\perp}$	Correlation times for molecular tumbling parallel and perpendicular to the symmetry axis (axially-symmetric case)
χ	(i) Magnetic susceptibility (footnote 7); (ii) nuclear quadrupole coupling constant ($\chi = e^2q_{zz}Q/h$)
$\omega_j, \omega_0, \omega_1, \omega_2$	As for $\nu_j, \nu_0, \nu_1, \nu_2$ but for angular frequencies.
Ω	Span of a tensor
Ω_1, Ω_2	RF magnetic fields, expressed in angular frequency units for a nucleus of magnetogyric ratio γ ($\Omega_1 = -\gamma B_1, \Omega_2 = -\gamma B_2$)

FOOTNOTES

- Note that confusion might arise when the so-called alphabet expansion is used for D , since this includes a term D which is not the dipolar coupling constant.
- M rather than m is frequently recommended, but most NMR practitioners use m so as to avoid confusion with magnetization.
- Whereas earlier IUPAC recommendations give a definition of δ which requires that the “unit” ppm is not stated when values are quoted, this is largely ignored and a change of recommendation is under consideration.
- $\zeta = 2\Delta\sigma / 3$.
- The symbols σ (and related terms for components), $\sigma_j, \sigma_{\parallel}, \sigma_{\perp}$ should refer to shielding on an absolute scale (for theoretical work). For shielding relative to a reference, symbols such as $\sigma_{\parallel} - \sigma_{\text{ref}}$ should be used.
- For tensors, doubled subscript *capital* letters X, Y and Z should generally be used for principal components, e.g. σ_{XX}, σ_{YY} and σ_{ZZ} for shielding. Alternatively, numerical subscripts may be used (e.g. $\sigma_{11}, \sigma_{22}, \sigma_{33}$).
- The symbol κ may also be used for magnetic susceptibility, some authors reserving χ for unrationalised units.

ACKNOWLEDGEMENTS

We are grateful to the following for helpful comments on the draft list: J.E. Bertie, J.W. Emsley, M. van der Graaf, H. Günther, K.-H. Hellwich, E. Hirota, H.-O. Kalinowski, C.L. Khetrepal, K. Kuchitsu, M.H. Levitt, J.C. Lindon, G.J. Martin, J. Mason, A.D. McNaught, P. Meadows, W.V. Metanomski, I.M. Mills, D.S. Moore, G.A. Morris, K.G. Orrell, A. Oskam, Z. Pajak, B. Petley, W. von Philipsborn, K. Salikhov, A.J. Thor, B.P. Winnewisser and K. Wüthrich.